



## Product Datasheet

<b>Product Name</b>	Clusterin Human Recombinant
<b>Cata No</b>	CB500180
<b>Source</b>	293 cell line (Human embryonic kidney).
<b>Synonyms</b>	CLI, AAG4, KUB1, SGP2, SGP-2, SP-40, TRPM2, MGC24903, Clusterin, Apolipoprotein J, Apo-J.

### Description

Clusterin also named Apolipoprotein J (APO-J) is a 75-80 kD disulfide-linked heterodimeric protein containing about 30% of N-linked carbohydrate rich in sialic acid but truncated forms targeted to the nucleus have also been identified.

The precursor polypeptide chain is cleaved proteolytically to remove the 22-mer secretory signal peptide and subsequently between residues 227/228 to generate the a and b chains. These are assembled in anti-parallel to give a heterodimeric molecule in which the cysteine-rich centers are linked by five disulfide bridges and are flanked by two predicted coiled-coil  $\alpha$ -helices and three predicted amphipathic  $\alpha$ -helices.

Across a broad range of species clusterin shows a high degree of sequence homology ranging from 70% to 80%. It is nearly ubiquitously expressed in most mammalian tissues and can be found in plasma, milk, urine, cerebrospinal fluid and semen. It is able to bind and form complexes with numerous partners such as immunoglobulins, lipids, heparin, bacteria, complement components, paraoxonase, beta amyloid, leptin and others. Clusterin has been ascribed a plethora of functions such as phagocyte recruitment, aggregation induction, complement attack prevention, apoptosis inhibition, membrane remodeling, lipid transport, hormone transport and/or scavenging, matrix metalloproteinase inhibition.

A genuine function of clusterin has not been defined.

One tempting hypothesis says that clusterin is an extracellular chaperone protecting cells from stress induced insults caused by degraded and misfolded protein precipitates.

Clusterin is up- or down regulated on the mRNA or protein level in many pathological and clinically relevant situations including cancer, organ regeneration, infection, Alzheimer disease, retinitis pigmentosa, myocardial infarction, renal tubular damage, autoimmunity and others.

The Clusterin Human contains a total 454 amino acids and a calculated molecular mass of 53058Da. The AA sequence (AA 1-427) is identical to Swiss-Prot-P10909 (AA 23-449, secreted Human Clusterin). C-terminal His tag 27AA (bold).

### Purity

Greater than 95% (SDS PAGE analyzed).

### Formulation

Filtered (0.4 micron) and lyophilized PBS, pH 7.5.

### Stability

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to **avoid repeated freezing/thawing cycles**. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

The lyophilized protein remains stable until the expiry date when stored at -20°C.

### Sequence

**\* For Non-Clinical Research Use Only \***



# California Bioscience

83103 Avenue 48, Ste.1B #204  
Coachella, CA 92236 USA  
Phone : +1.6268339877  
Email : [info@cali-bio.com](mailto:info@cali-bio.com)

DQTVSDNELQ EMSNQGSKYV NKEIQNAVNG  
VKQIKTLIEK TNEERKTLLS NLEEAKKKKE  
DALNETRESE TKLKELPGVC NETMMALWEE  
CKPCLKQTCM KFYARVCRSGS GLVGRQLEE  
FLNQSSPFYF WMNGDRIDSL LENDRQQTHM  
LDVMQDHFSRA SSIIDELFQ DRFFTREPQD  
TYHYLPFSLP HRRPHFFFPK SRIVRSLMPF  
SPYEPLNFHA MFQPFLEMIH EAQQAMDIHF

HSPAFQHPPT EFIREG **Product Datasheet**  
TGCLRMKDQC DKCREILSVD  
CSTNNPSQAKLRREDESQ VAERLTRKYN  
ELLKSYQWKM LNTSSLLEQL NEQFNWVSRL  
ANLTQGEDQYYLRVTTVASH TSDSDVPSGV  
TEVVVKLFDS DPITVTVPE VSRKNPKFME  
TVAEKALQEY RKKHREE **GSL GSGWSHPQFE**  
**KTGHHHHHHH HGGQ**

**\* For Non-Clinical Research Use Only \***